

UGC Journal Details

Name of the Journal : Journal of Emerging Technologies and Innovative Research

ISSN Number : 23495162

e-ISSN Number :

Source: UNIV

Subject: Electrical and Electronic Engineering


Publisher: JPUBLICATION

Country of Publication: India

Broad Subject Category: Science

{ Print




Co-ordinator
I.Q.A.C.
KVN Naik Arts, Commerce
& Science College, Canada Corner,
Nashik-422 002.

Impact of Endosulfan on Neurosecretory cells of Land Slug, *Laevicaulisalte*.

D.B.Goswami

K.V.N.Naik Arts, Commerce and Science college P.G Department of Zoology, Nashik, Maharashtra.

ABSTRACT: Incorporation of the toxic compound or their metabolites in lower organisms, as well as in the tissues of fishes, birds and humans have been recorded to cause serious morphological alterations in vital tissues of organisms even at very low levels. So the studies on the effects of pesticides on molluscan life are an important aspect of pollution. The studies on effect of the lethal concentration (LC_{50}) of pesticide endosulfan on the neurosecretory cells of the central ganglia such as cerebral, visceral and pedal ganglia have been carried out during pre-reproductive of land slug, *Laevicaulisalte*. Method- Slugs were treated with lethal concentration (LC_{50} concentration) of pesticide, endosulfan for 96 hr period during pre-reproductive. Sections of central ganglia (cerebral, visceral and pedal) were stained with Mallory's triple, Gomery's chrome-hematoxylin phloxin (CHP) staining method. All the observations were made under light microscope to evaluate the changes at cellular level due to pesticide, endosulfan and photomicrographed.

Key words: endosulfan, LC_{50} , *Laevicaulisalte*.

Introduction:
In mollusca, neurosecretory cells were first discovered by Scharrer (1935) in the cerebral and visceral ganglia of *Aplysia limacina* and *Leurobranchaea*. Perusal of literature reveals that the aspect of neurosecretion in molluscs has been reviewed by Gabe (1965, 1966), Lubet (1966, 1973), Martoja (1972) and Golding (1974). The development of the subject has been hampered by the presence of shell, very small ganglia in small snails, diffused distribution of NSCs, and by the ignorance of the chemical nature of the neurohormones. In gastropods, the nervous system is constructed on the same general plan as in the other molluscs undergo. (Ahirrao et al., 2012) Amongst invertebrates, molluscs show great variability in their nervous system, ranging from primitive arrangement in chitons to the complex mass of fused ganglia forming the "brain" of cephalopods. Pharmacological and physiological aspects of molluscs using effector organs have received considerable attention (Huadart, 1975; Bayne, 1976). Evidence for the occurrence of a wide variety of neurotransmitters in different tissues of bivalve molluscs including the cerebral ganglia has been discussed from the functional point of view by Leak and Walker (1980). Gabe (1955) reported the presence of neurosecretory cells in 20 marine lamellibranch molluscs. Lubet (1955b) described the relationship between neurosecretion and sexual cycle in *Stilus* and *Chlamys*. Thus, there are relatively very few reports on the impact of pesticides on the histopathological changes in molluscs. Most of the work in this connection has been dealt in vertebrates, especially on the fishes. Since little work is carried out on slugs, except on neurosecretion and reproductive system of *L. alte* (Bodhankar, 1984; Vyawahare, 1988; and Jawalikar, 1989). In the present study an attempt has been made to study the effect of one of the tested pesticides, endosulfan on the neurosecretory cells of the central ganglia such as cerebral, visceral and pedal ganglia have been carried out during pre-reproductive, reproductive and post reproductive period of land slug, *Laevicaulisalte*.

MATERIAL AND METHODS

Healthy specimens of *Laevicaulisalte* were collected from the cultivated fields in and around Hinganwadi, Pale and Kalwan area of Nashik and maintained under laboratory conditions in troughs with sufficient amount of moist soil. They were fed once in a day on carrot, beetroot or calotropis leaves. They were kept in laboratory for three days before subjected to experimentations. Healthy and mature animals of equal size were selected and were starved during experimental period. Slugs were treated with lethal concentration (LC_{50}) of pesticide, endosulfan for 96 h during pre-reproductive. After 96 h acute toxicity of endosulfan, central ganglia (cerebral, visceral and pedal) belonging to control and experimental groups (LC_{50} concentrations) were removed and fixed in aqueous Bouin's fluid fixative. The tissues were then dehydrated in ethyl alcohol, cleared in xylene and embedded in paraffin wax (58-60°C). The sections were cut at 6-7 μ m thickness on a rotary microtome. Sections of central ganglia (cerebral, visceral and pedal) were stained with Mallory's triple, Gomery's chrome-hematoxylin phloxin (CHP) staining method. All the observations were made under light microscope to evaluate the changes at cellular level due to pesticide, endosulfan and photomicrographed.

RESULTS:




Fig. -(1-A-B-)Changes in the neurosecretory cells from cerebral ganglia of *Laevecaulisalte* due to pesticides, Endosulfan during pre - reproductive period.
A - control ; B - Experimental group (x 100)
(a - pyriform cells; ax - axon; b- oval cells; n - nucleus; nu - nucleolus;
nsm - neurosecretory material)

The Neurosecretory cells from cerebral ganglia :

The observations of the cerebral, visceral and pedal ganglia showed the presence of cells, which are cytologically different and longer than ordinary, numerous neuronal cells. The neurosecretory cells are distributed along the periphery of the ganglia and each possesses a large nucleus with prominent cytoplasm. Their axons and perikaria are more or less filled with the secretory material stained conspicuously by different seasons.

Control : (fig- 1-A) Based on the differences in their size, shape, presence or absence of vacuoles and staining properties of secretory material, the two types of neurosecretory cells could be identified, cell type 'a' or pyriform cells and cell type 'b' or oval cells.

The changes in the neurosecretory material from cerebral, visceral and pedal ganglia was as follows:
Cerebral: (fig- 1-B) The secretory material in the cytoplasm of both the types of cells was granular. Most of the cells of type a contained large amount of secretory material while others contained no granules. The secretory material in the hillock and/or in the axons. Vacuoles in the cytoplasm appeared in large size in a few cells due to release of secretory material. In cell type 'b' the vacuoles are large but showed no characteristics shape.




Fig. (2-A-B) Changes in the neurosecretory cells from visceral ganglia of *Laevecaulisalte* to pesticides, Endosulfan during pre - reproductive period.
control ; B - Experimental group (x 100)
pyriform cells; ax - axon; b- oval cells; n - nucleus; nu - nucleolus;
nsm - neurosecretory material)

Cerebral : The neurosecretory material accumulated along the periphery of the nuclei, being more in type 'a' cells. Nuclei and nucleoli stained prominently. In type 'b' cells the secretory material was scattered in the cytoplasm in many cells.




Fig. (3-A-B) Changes in the neurosecretory cells from pedal ganglia of *Laevicaulisalte* due to pesticides, Endosulfan during pre-reproductive period.

A - control; B - Experimental group (x 100)

a - axon cells; ax - axon; b - oval cells; n - nucleus; nu - nucleolus; nsm - neurosecretory material; p - perikaryon

Observation: The neurosecretory material was less vacuolated, however, traces of material were recorded around the nuclei in both the types of cells (Fig. 3-A).

REFERENCE:

- [1] Stronganova, N.S. (1963): Peculiarities of spermatogenesis in certain bivalve molluscs. *VestnMosk. Univ. Ser. Biol. Soil. Sci.*, 6: 25-34.
- [2] Umiji, S. (1969). Neurosecretion in *Mytilus perna* (Mollusca Lamellibranchia). *Zool. Biol. Mar. (Sao Paulo) Nova. Ser.*, 26: 181-254.
- [3] Bayne, B. L. (1976). *Marine mussels: Their ecology and physiology*, Cambridge University Press, Cambridge, London, New York, Melbourne, 1-495.
- [4] Srivastava, V.M.S. and R.S. Tripathi (1981): Histopathological changes in the tissue of *Esomus dandricus* (Ham.) under sodium and ammonium sulphate stress. *Comp. Physiol. Ecol.*, 6(4): 236-238.
- [5] Akarte, S.R. (1985). Effect of organophosphorus insecticides on bivalve molluscs. Ph.D. Thesis, Marathwada University, Aurangabad, M.S., India, 1-190.
- [6] Shrivastava, R. S.; L. K. Mudgal; G. D. Sharma and D. K. Verma (1987): Histopathological study and Gas Chromatographical Analysis of Pesticide Residue (Thiodon) on the Gills of *Heteropneustes fossilis*. *Proc. of the Natl. Symp. on Environ. Poll. And Pesticide Toxicol.* Held at University of Jammu, Jammu (J & K) from Dec. 10.12.1987 pp. 31-37.
- [7] Wester, P.W. (1991): Histopathological effects of environmental pollutants B-HCH and methyl mercury on reproductive organs in freshwater fish. *Comp. Biochem. Physiol.* 110C (1/2): 237-239.
- [8] Magare SR. 1991. Endocrine regulations in a terrestrial snail, *Cerastus moussonianus*. Ph. D. Thesis, Dr. BAMU, Aurangabad
- [9] Patil P.N, Rao K.R, Sasane S.R, Vedpathak A.N, Shejule K.S and Chaudhari T.R. (1993) Impact of Zolone on the organic constituents of the Freshwater gastropod, *Thiara lineata* from Panzara river at Dhule. *Proc. Acad. Environ. Biol.*, 2(2):193-197.
- [10] Patil VN and Kulkarni AB. (2000) Effect of heavy metals on neurosecretory cells of a freshwater Prosobranch snail, *Ballamya bengaliensis*. *Environmental Issues and Sustainable Development* Vinit Publications, Aurangabad. 10-12.
- [11] Patil S.S. and Mane U.H. (2001). Mercury impact on the bivalve *L. marginalis* in summer season: biochemical and histological approach. *Environmental and Agriculture: Biodiversity, Agriculture and pollution in South Asia*, pp. 581-587.
- [12] Ahirrao KD1, Kedkar GD2. (2012) Variations in LC50 of some pesticides due to physico-chemical parameters and impact of sevin on neurosecretory cells of a fresh water snail, *Thiara lineata*. *Journal of Research in Biology*: 050-056
- [13] Vijaya Kumar Kanapala and A.S. Priyadarsini Arasada, 2015. Histology and Cytochemistry of the Neurosecretory Cells (NSC) of the Freshwater Snail *Lymnaea luteola* (Lamarck) Mollusca: Gastropoda. *International Journal of Zoological Research*, 11: 215-221.