# **UGC Journal Details**

	Name of the Journal :	Research Journey
	ISSN Number :	
en e	e-ISSN Number :	23487143
	Source:	UNIV
	Subject:	Marathi
	Publisher:	Swatidhan International Publication
	Country of Publication:	India
	Broad Subject Category:	Multidisciplinary

Print



Co-ordinato I.Q.A.C. KVN Naik Arts, Commerce & Science College, Canada Corner, Nashik-422 002.





# विश्वविद्यालय अनुदान आयोग University Grants Commission

quality higher education for all

(https://www.ugc.ac.in/faq.aspx) RTI (https://www.ugc.ac.in/subpage/RTI-Act.aspx) Tenders (https://www.ugc.ac.in/Tenders.aspx) Jobs (https://www.ugc.ac.in/ugc\_jobs.aspx) Press Release (https://www.ugc.ac.in/ugc\_pressrelease.aspx)

Contact Us (https://www.ugc.ac.in/contact.aspx)

# UGC Approved List of Journals

nttps://www.ugc.ac.in/hindi/)

|| Home (journal\_list.aspx)

You searched for r

### Total Journals : 1436

how 25	50	▼ entrie	sSearch: Rese	arch journey		
Nrta, Commerce Hege, Cenada Cornely hik-422 002. weiV	SI.No.	Journal No	Title	Publisher	ISSN	E- ISSN
View (ugc_admin_journal_report.aspx? eid=NDA3MDU=)	1168	40705	Research Journey International E Research Journal	Swatidhan Pub. Yevla		23487143
View (ugc_admin_journal_report.aspx? eid=NDQxMTc=)	1220	44117	Research Journey	Swatidhan International Publication		23487143

### For UGC Officials

> e-Office (https://ugc.eoffice.gov.in/acl\_users/credentials\_cookie\_auth/require\_login? came\_from=https%3A//ugc.eoffice.gov.in/)

- > NIC mail (https://mail.gov.in/iwc\_static/c11n/allDomain/layout/login\_gov2.jsp?
- lang=en&3.0.1.2.0\_15121607&svcs=abs,im,mail,smime,calendar,c11n)
- -RTI online (https://rtionline.gov.in/RTIMIS/login/index.php)
- > PG portal (https://pgportal.gov.in)

nttps://www.ugc.ac.in/journallist/subjectwisejurnallist.aspx?tid=cg==





# Theoretical and Infrared studies of Ethyl methyl ketone, Methyl isobutyl ketone, Propiophenone and Cyclohexane with o-dichlorobenzene and ochlorophenol

### Dr. V. J. Naukudkar

Head, Department of Chemistry, K. V. N. Naik, Arts, Commerce & Science College, Nashik, 422 002,

### Abstract:

Binary mixtures of o-dichlorobenzene and o-chlorophenolwithEthyl methyl ketone, Methyl isobutyl ketone, Propiophenone and cyclohexane have been prepared at room temperature for entire concentrations. These binary mixtures studied by Infrared spectroscopy. The data have been utilized to compute the Infrared studies correlate to Theoretical Calculations. The results have been interpreted in terms of molecular interactions existing between the components of the mixtures.

Keywords: Infrared spectroscopy, Theoretical studies, DFT/B3LYP method, 6-31G (d) basis set, o-Dichlorobenzene, o-Chlorophenol, Ketones, Cyclohexane, Molecular Interactions.

#### 1. Introduction

The composition dependence of thermodynamic properties has proved to be very useful tools in understanding the nature and extent of pattern of molecular aggregation resulting from intermolecular interaction between components<sup>1, 2</sup>. In the present paper binary liquid mixture of o-dichlorobenzene (ODB) and o-chlorophenol (OCP) with Ethyl methyl ketone (EMK), Methyl isobutyl ketone (MIBK), Propiophenone (PN) and cyclohexane (CYN)have been studied. Theoretical and Infrared Red studiesof odichlorobenzene (ODB) and o-chlorophenol (OCP) with above ketones have been studied over the entire composition range.

#### 2. Experimental

The entire chemical used was of A. R. grade with purity >99% are used as such.Binary mixtures were prepared by mixing a known mass of each liquid in an airtight, stoppered glass bottle.I.R. measurement through liquid mixtures provides an excellent tool to investigate inter and intermolecularinteractions between like and unlike molecules.I.R. Spectroscopy studies of pure and binary mixtures of ODB and OCP with ketones have been recorded over the whole composition range at room temperature and atmospheric pressure. FTIR spectra were recorded on a FTIR spectrometer (Model: SIMADZU 8400S PC) by using KBr pellet in the region 400-4000cm<sup>-1</sup> with 4.0 cm<sup>-1</sup> resolution. The transmission values were read in steps of 5%. The spectrometer possesses out to aligned energy optimization and dynamically aligned interferometer. It is fitted with KBr beam splitter, a DLATGS detector. A base line correction was made for the spectra recorded. The data have been utilized to compute the Infrared studies correlate to Theoretical Calculations.

#### 3. Theory

An IR is widely used for: (1) identification and determination of structure (2) determining the purity and quantitative analysis (3) following the course of reaction (4) hydrogen bonding (5) molecular geometry and conformational analysis (6) chemistry of organic polymers and (7) reactions of reactive species like free radicals and ions.

The stretching frequency of chemical bond depends upon (i) bond strength (ii) reduced masses of the atoms forming bond. Any factor which will increase bond strength will increase stretching frequency of the bond, and if the mass of atoms forming the chemical bond is increased, the reduced mass will increase and stretching frequency will decrease. The following factors affect the IR absorption frequency: (a) Inductive and resonance effect (b) Conjugation (c) size of the ring and (d) hydrogen bonding.

The free (non-bonded) -OH group is observed only in very dilute solution in non-polar solvent. The free O-H stretching band appears at about 3600cm<sup>-1</sup>. Intermolecular H-bonding increases with the increase in concentration of solution. A broad O-H stretching vibration is obtained for intermolecular hydrogen bonding in the range from 3400-3200cm<sup>-1</sup>. In short, H-bonding decreases the O-H stretch frequency. Intermolecular hydrogen bonding weakens the O-H bond, thereby shifting the band to lower frequency.

#### 4. **Results and Discussion**

In recent years, experimental results have been frequently supported with spectroscopic/theoretical calculations which have proved quite helpful in explaining the comparative relativities and many other properties of the organic molecules/complexes.Hydrogen bonding is of central importance in the molecular sciences for both practical and theoretical reasons<sup>3-10</sup>. Hydrogen bonding has held particular interest in recent

	RESEARCH JOURNEY International Multidisciplinary E-R	ISSN- 2348-7143	
CHRONICUMMET	Impact Factor - (SJIF) – 6.261, (CIF) - 3.452, (GIF) –0.676 Special Issue - 118 Modern Trends in Green Chemistry, Medicinal Chemistry and environmental Science	February 2019	UGC Approved No. 40705

years due to the central role it plays with regard to molecular recognition in both biological and artificial systems<sup>11-14</sup>.

There has been increasing interest in C-H--O hydrogen bonds recently, both theoretically and experimentally<sup>15-17</sup>. Because carbon is not particularly electronegative, the ability of the C-H group to serve as a proton donor in hydrogen bonding depends on the carbon hybridization [C (sp)-H > C (sp<sup>2</sup>)-H > C (sp<sup>3</sup>)-H] and on the electron with-drawing strength of adjacent substituent<sup>18-20</sup>. Although the C-H--O interaction energy varies and is much weaker than conventional hydrogen bonds, it can play important role in molecular conformation<sup>21</sup>, crystal packing<sup>22</sup> and protein folding<sup>23</sup>. Close C-H--O contracts have been observed widely in high resolution protein structures.Extensive calculations are carried out to determine the optimized geometries, and electronic structure and thermochemical data.On the basis of semi-empirical calculations, the reactivity of heterocyclic compounds towards electrophilic substitution has been predicted<sup>24-29</sup>.

IK spectroscopy is the leading method for identification of hydrogen bonds. For instance the formation of an O-H--O hydrogen bond elongates and weakens the O-H bond. The resulting red shift of the O-H bond stretching frequency can be easily detected in the IR spectra and its magnitude indicates the strength of the hydrogen bond<sup>30</sup>. This phenomenon is called anti-hydrogen bond<sup>31-32</sup>. However, Scheineret al.<sup>33</sup> have concluded, from a set of careful calculations that "antihydrogen" bonds do not differ fundamentally from conventional hydrogen bonds. According to their results, the electron density redistribution upon hydrogen bond formation is similar for both the C-H--O and O-H--O interactions. The amount of charge transferred from a donor to an acceptor is roughly proportional to the bonding strength. The solvent structure determines the nature of molecular interaction. Spectral changes are caused by hydrogen bonding, van der Waals interactions, dipole-dipole interactions, induced-dipole interactions etc. Information about hydrogen bonding is derived from the bond width, position and intensity of the first overtone band soft the -OH band (intensity of first overtone band decreases)<sup>34</sup>.

Spectrum of some OCP + ketones show two absorption bands -OH. The sharp band is due to free -OH while weak band is due to hydrogen bonded -OH. Presence of hydrogen bonding between -OH of OCP with oxygen of studied ketones and absence of hydrogen bonding in systems of ODB + ketones is confirmed by FTIR. In case of ODB + Ketones, carbonyl stretching frequency is not so affected but slightly increases at  $x_1 \approx 0.8$  and 0.9.

The interesting behavior of intramolecular hydrogen bonds both from the practical and theoretical points of view has received increased attention in recent years<sup>35-43</sup>. This can be attributed to the fact that the value of hydrogen bond energy is only a few kilocalories per mole and can readily perturb by any change in its environment. This is especially important in studies of weekly bound complexes of ketones and OCP.

### 4.1 Infrared studies

RE

The FTIR absorption frequencies of the binary mixtures of ODB and OCP with ketones over the entire range at room temperature are listed in Tables 1.0-3.0.

### 4.1.1 ODB + Ketones

C=O stretching vibrations: The band due to C=O stretching vibration is observed in the region 1650-1850 cm<sup>-1</sup>. For aliphatic ketone C=O stretching vibration is at 1710-1720 cm<sup>-1</sup> and for aromatic ketone C=O stretching vibration it is observed at around 1680-1690 cm<sup>-1</sup> due to conjugation effect. From Table 1.0, it is observed that C=O stretching frequency is almost same from  $x_1 \approx 0.1$ - 0.9 for ODB + EMK and ODB + MIBK. Lower C=O stretching frequency is observed at  $x_1 \approx 0.1$  and higher C=O stretching frequency is observed at  $x_1 \approx 0.1$  and higher C=O stretching frequency is observed at  $x_1 \approx 0.1$  and higher C=O stretching frequency is observed at  $x_1 \approx 0.1$  and higher C=O stretching frequency is observed at  $x_1 \approx 0.1$  and higher C=O stretching frequency is observed at  $x_1 \approx 0.1$  and higher C=O stretching frequency is observed at  $x_1 \approx 0.1$  and higher C=O stretching frequency is observed at  $x_1 \approx 0.1$  and higher C=O stretching frequency is observed at  $x_1 \approx 0.1$  and higher C=O stretching frequency is observed at  $x_1 \approx 0.1$  and higher C=O stretching frequency is observed at  $x_1 \approx 0.1$  and higher C=O stretching frequency is observed at  $x_1 \approx 0.8$  and 0.9 for all systems except ODB + PN. The frequency follows the order: ODB + MIBK > ODB + EMK > ODB + PN.

X <sub>1</sub>	ODB+EMK	ODB +MIBK	ODB +PN
0	1716.5(s)	1716.6(s)	1685.7(s)
0.1	1716.5(w)	1716.7(s)	1685.8(s)
0.2	1716.5(w)	1716.7(s)	1687.7(s)
0.3	1716.5(s)	1716.7(s)	1687.7(s)
0.4	1716.5(m)	1716.7(s)	1687.7(s)
0.5	1716.5(s)	1716.7(s)	1687.7(s)
0.6	1716.5(s)	1716.7(s)	1687.7(s)

### Table 1. IR stretching frequencies of C=O (cm<sup>-1</sup>) in $(x_1)$ ODB and $(1-x_1)$ ketones systems.



## RESEARCH JOURNEY International Multidisciplinary E-Research Journal

Impact Factor - (SJIF) – 6.261, (CIF) - 3.452, (GIF) –0.676 Special Issue - 118 February 2019 Modern Trends in Green Chemistry, Medicinal Chemistry and environmental Science

UGC Approved No. 40705

1.0			
0.9	1716.5(m)	1716.7(m)	1689.7(m)
0.8	1714.6(s)	1716.7(m)	1689.7(m)
0.7	1716.5(s)	1716.7(m)	1687.7(s)

w = weak absorption, s = strong absorption, m = medium absorption.

### 4.1.20CP + Ketones

C=O stretching vibrations: From Table 2.0, it is observed that C=O stretching frequency decreases from  $x_1 \approx 0.1$  to 0.9 for OCP + EMK, OCP + PN and OCP + MIBK. Higher C=O stretching frequency observed at  $x_1 \approx 0.1$ . The frequency follows the order: OCP + EMK > OCP + MIBK > OCP + PN. Table 2. IR stretching frequencies of C=O (cm<sup>-1</sup>) in ( $x_1$ ) OCP and (1- $x_2$ ) ketones systems.

<b>X</b> <sub>1</sub>	OCP+EMK	OCP+MIBK	OCP+PN
0	1716.5(s)	1714.6(s)	1685.7(s)
0.1	1703.0(s)	1716.7(s)	1685.8(s)
0.2	1703.0(s)	1707.0(s)	1685.8(s)
0.3	1703.0(s)	1705.1(s)	1685.8(s)
0.4	1703.0(s)	1701.2(s)	1681.9(s)
0.5	1703.0(s)	1701.2(s)	1674.2(s)
0.6	1701.1(s)	1697.4(m)	1674.2(s)
0.7	1701.1(s)	1697.4(m)	1672.3(s)
0.8	1699.2(s)	1697.4(m)	1672.3(s)
0.9	1697.2(s)	1697.4(w)	1670.4(s)
1.0			

**O-H stretching vibrations:** The non hydrogen-bonded hydroxyl group of phenol absorbs strongly in the 3584-3650 cm<sup>-1</sup> region. The sharp free hydroxyl bands are observed in the vapour phase and in dilute solution in non polar solvent<sup>44</sup>. Intermolecular hydrogen bonding increases as the concentration of solution increases and additional bands appears at lower frequencies 3200-3500 cm<sup>-1</sup>. Hence in the present investigations (Table 3.0); phenolic O-H frequency is low at  $x_1 \approx 0.5$  and decreases from  $x_1 \approx 0.1 - 0.5$  and again increases onwards for OCP + EMK. This frequency increases from  $x_1 \approx 0.1 - 0.9$  for OCP + PN. IR frequencies of phenolic O-H remains same in  $x_1 \approx 0.2 - 0.5$  and increases from  $x_1 \approx 0.6 - 0.9$  for OCP + MIBK. For OCP systems the sharp band is due to free -OH, while weak band is due to hydrogen bonded - OH. Two bands are observed in OCP + PN, OCP + MIBK in  $x_1 \approx 0.6 - 0.9$  and one sharp band is recorded for OCP + EMK over entire range of composition.

## Table 3. JR stretching frequencies of O-H (cm<sup>-1</sup>) in (x<sub>1</sub>) OCP and (1-x<sub>1</sub>) ketones systems.

<b>X</b> <sub>1</sub>	OCP+ EMK	<b>OCP+ MIBK</b>	OCP+ PN
0	,		
0.1	3402.2(s)	3352.3(w)	3360.1(w)
0.2	3394.5(s)	3296.7(s) 3292.6(s)	3360.1(s)
0.3	3400.3(s)	3298.3(m) 3392.9(s)	3392.9(s) 3365.9(s)
0.4	3396.4(s)	3296.4(Broad & s) 3524.0(w)	3390.9(s) 3369.7(m)
0.5	3390.6(s)	3398.6(Broad & s) 3524.0(w)	3396.7(s)
0.6	3404.1(s)	3392.9(Broad & s) 3524.0(w)	3404.4(s) 3524.0(w)
0.7	3425.3(s)	3521.8(Broad & s) 3524.0(m)	3416.0(s) 3524.0(m)

	RESEAR	ISSN- 2348-7143			
SCARDINGUNKET	Impact Factor Modern T	cience February 2019	UGC Approved No. 40705		
	0.8	3431.1(s)	3524.0(s) 3423 7(m)	3421.8(s) 3524.0(s)	

0.8		3423.7(m)	3524.0(s)
	3446.6(s)	3524.0(s)	3448.8(m)
0.9		3458.4(w)	3524.0(s)
1.0	3456.2(s)	3456.2(s)	3456.2(s)

The position and breadth of an -OH absorption band depends on the concentration of the solution. The more concentrated solution and the more -OH containing molecules form intermolecular hydrogen bonds. It is easier to stretch an O-H bond if it is hydrogen bonded, because the hydrogen is attached to oxygen of a neighboring molecule. Therefore, the stretch of a concentrated (hydrogen bonded) solution of OCP occurs at 3427.3 cm<sup>-1</sup>. However no such dilution effect is observed in other binary mixture. But other all systems show hydrogen bond effect at higher concentration ( $x_1 \approx 0.1$  to 0.6) of OCP. Hydrogen bonded O-H group have broader absorption bands (e.g. OCP + MIBK; Table 3) while that of non-hydrogen-bonded O-H groups are shaper (e.g. OCP + EMK; Table 3).



It is seen that the O-H stretching frequency increases when mole fraction of OCP decreases. Higher values of O-H stretching frequency is observed at high mole fraction and lower values of O-H stretching frequency is recorded in between 0.2 to 0.5 mole fractions of OCP. This may be due to hydrogen bonding between phenolic O-H group and carbonyl group. The O-H stretching frequency of OCP is 3456.2 cm<sup>-1</sup>, which gets decrease in presence of ketone because of intermolecular hydrogen bonding between them. The decrease in -OH stretching frequency is not same in all compositions of OCP + ketone systems. The binaries OCP + PN and OCP + MIBK show large decrease in the O-H stretching frequency, but in other binary systems OCP + EMK, the decrease is small.



Spectrum of some OCP + ketones show two absorption bands -OH. The sharp band is due to free -OH while weak band is due to hydrogen bonded -OH. Presence of hydrogen bonding between -OH of OCP with oxygen of studied ketones and absence of hydrogen bonding in systems of ODB + ketones is confirmed by FTIR. In case of ODB + Ketones, carbonyl stretching frequency is not so affected.

The interesting behavior of intramolecular hydrogen bonds both from the practical and theoretical points of view has received increased attention in recent years<sup>35-43</sup>. This can be attributed to the fact that the value of hydrogen bond energy is only a few kilocalories per mole and can readily perturb by any change in its environment. This is especially important in studies of weekly bound complexes of ketones and OCP.

	RESEARCH JOURNEY International Multidisciplinary E-R	ISSN- 2348-7143	
RESEARCHULUPHEY	Impact Factor - (SJIF) – 6.261, (CIF) - <b>3.452</b> , (GIF) – <b>0.676</b> Special Issue - 118 Modern Trends in Green Chemistry, Medicinal Chemistry and environmental Science	February 2019	UGC Approved No. 40705

Fig 4.1.16 IR streaching frequencies of O - H (cm<sup>-1</sup>) for OCP  $\div$ PN system.



### 4.2 THEORETICAL STUDIES

All the theoretical calculations on OCP/ODB and their mixtures with ketones were performed by the use of G03W series of programs<sup>45</sup>. Geometries of optimizations for all the investigated molecules in this work were carried out using the DFT/B3LYP method with medium size 6-31G (d) basis set.

The difference between the observed and scaled frequency values of most of the fundamentals in many cases is very small. Any discrepancy noted between the observed and calculated frequencies may be due to the fact that the calculations have been actually done on the single molecule/complex between two molecules in the gaseous state contrary to the experimental values recorded in the presence of intermolecular interactions. Therefore the assignment made at higher levels of theory with only reasonable deviations from the experimental values seem to be correct.

#### 4.2.1 Hydrogen bond (Table 4.2.1)

Maximum hydrogen bond distance is noted for Cis-OCP + PN and minimum distance (i.e. strongest hydrogen bond among complexes) for Trans-OCP + MIBK. Weak intermolecular hydrogen bonds are also seen in Cis-OCP + EMK. Strong intermolecular hydrogen bonds are also observed for Trans-OCP + EMK. Table 4.2.1:Optimized atomic distances of systems at B3LYP level using 6-31G (d) basis set.

Sr. No.	System	О-Н	C-Cl	C=0	=C-0	ОН	Cl0=C
1.	ODB		1.81381				
2.	Trans-OCP+EMK	0.98524	1.75649	1.22645	1.35269	1.80637	5.69171
3.	Trans-OCP+PN	0.98239	1.75656	1.23165	1.35365	1.80984	5.70860
4.	Trans-OCP+MIBK	0.98466	1.75684	1.22638	1.35268	1.79973	5.71677

C

		RESEARCH JO	DURNEY	Internation	al Multidisc	iplinary E-R	esearch Journa	ISSN- 2348-7143
SEAFGHOUGHMET		Impact Factor - (SJIF) – 6.261, (CIF) - 3.452, (GIF) –0.676 Special Issue - 118 Modern Trends in Green Chemistry, Medicinal Chemistry and environmental Science			February 201	9 UGC Approved No. 40705		
	5.	Trans-OCP+CYN	0.96954	1.75164		1.36207		
	6.	C.s- OCP+EMK	0.98448	1.76171	1.22355	1.35122	1.88485	3.34944
	7.	Cis-OCP+PN	0.97402	1.76255	1.22350	1.35693	4.20730	3.39307
	8.	Cis-OCP+MIBK	0.98490	1.76219	1.22466	1.35095	1.87845	3.36838
	9.	Cis-OCP+CYN						
	10.	ODB+EMK	-	1.74471 1.75198	1.21824			3.01114 5.46307
	11.	ODB+PN	-	1.74429 1.75157	1.22459	-	-	3.12476 5.52668
	12.	ODB+MIBK		1.75159 1.74502	1.21920			3.10924 5.51199
	13.	ODB+CYN		1.74854 1.74824	-			

Hydrogen bond: O(C=O of KETONE) -- H (OH of OCP)-

4.2.2 Atomic Charges (Table 4.2.2)

Atomic charges unlike the electron density are not a quantum mechanical observable and are not unambiguously practicable. Gaussian jobs perform a Mulliken population analysis, which partitions the total charge among the atoms in the molecule/complex. Mulliken population analysis is an arbitrary scheme for assigning changes. Both chlorine atom of free ODB shows positive charge. In present investigation (Table 4.2.5), for ODB + EMK, ODB + PN and ODB + MIBK system only one of the chlorine atom shows positive charge and other shows negative charge. In ODB + CYN both chlorine atom of ODB show positive value but lower than that of free ODB.

In case of ketones, negative charge is estimated on oxygen of C=O. It is maximum negative charge for AP and minimum for AC. For other complexes, it is maximum negative for Trans-OCP + PN. maximum negative charge is noted on oxygen of -OH for complexes Cis-OCP + EMK and Cis-OCP + MIBK and minimum for Cis-OCP + PN. In all complex system, Trans-OCP + CYN shows lowest negative charge on oxygen of -OH. Positive charge on hydrogen atom of -OH decreases in Cis-OCP + PN. In all complex system, Trans-OCP + CYN shows lowest value positive charge on hydrogen atom of -OH.

Chlorine of OCP shows negative charge for all Cis/Trans-OCP + Ketone systems in present investigation (Table 4.2.5). It is observed that minimum for Cis-OCP + PN. In OCP + CYN, chlorine atom of OCP shows lowest negative value. In case of ketones, positive charge on carbon of C=O, is maximum for EMK. While for complexes; it is maximum for Trans-OCP + EMK and Trans-OCP + MIBK and minimum for Trans-OCP + PN and Cis-OCP + PN.

Table 4.2.2: Optimized Mulliken atomic charges of systems at B3LYP levelusing 6-31G (d) basis set.

S r	Committee Latif	1 ( A		0.1410
. System N o	<c-o-h (degree)</c-o-h 	<cl-c=c (degree)</cl-c=c 	<o-c=c (degree)</o-c=c 	<c-c=o (degree)</c-c=o 
1 ODB	-	121.544		
2 PN	-		20 <b>-</b> 1	119.780 120.303°
3 MIBK	-	_		121.496 121.632°

23

F	RESEARCH JOU	RNEY Interna	ational Multidis	ciplinary E-Re	esearch Journal	ISSN- 2348-714
Ir	mpact Factor - (SJIF) – 6.261, Modern Trends in Green Cher	(CIF) - 3.452, (GIF) nistry, Medicinal Chem	-0.676 Special istry and environment	Issue - 118 al Science	February 2019	No. 40705
4	CYN			-		
5	Trans-OCP+EMK	109.956	119.319 <sup>a</sup> 119.633 <sup>b</sup>	121.579 <sup>c</sup> 120.125 <sup>d</sup>	118.913 122.879°	
6	Trans-OCP+PN	109.956	119.348 <sup>a</sup> 119.633 <sup>b</sup>	122.859 <sup>c</sup> 118.846 <sup>d</sup>	118.235 121.443°	
7	Trans-OCP+MIBK	109.952	119.316 <sup>a</sup> 119.633 <sup>b</sup>	122.846 <sup>c</sup> 118.921 <sup>d</sup>	120.476 121.305°	
8	Trans-OCP+CYN	108.911	119.540 <sup>a</sup> 119.923 <sup>b</sup>	122.735 <sup>c</sup> 118.491 <sup>d</sup>		
9	Cis-OCP+EMK	114.070	119.331 <sup>a</sup> 119.370 <sup>b</sup>	125.055° 117.150 <sup>d</sup>	120.174 121.737°	
1 0	Cis-OCP+PN	109.838	120.162 <sup>a</sup> 118.617 <sup>b</sup>	123.703° 117.909 <sup>d</sup>	119.930 120.156 <sup>e</sup>	-
1 1	Cis-OCP+MIBK	114.120	119.289 <sup>a</sup> 119.389 <sup>b</sup>	125.087° 117.147 <sup>d</sup>	120.598 121.809°	
1 2	Cis-OCP+CYN	109.508	$118.587^{a}$ $119.990^{b}$	123.652 <sup>c</sup> 118.003 <sup>d</sup>		
1 3	ODB+EMK		121.330 <sup>a</sup> 121.809 <sup>b</sup>	-	120.903 121.627°	
1 4	ODB+PN	-	121.338 <sup>a</sup> 121.849 <sup>b</sup>	-	119.937 120.075°	
1 5	ODB+MIBK		121.339 <sup>a</sup> 121.836 <sup>b</sup>		121.191 121.863°	
1 6	ODB+CYN	-	121.499 <sup>a</sup> 121.503 <sup>b</sup>	-		

### Energy and dipole moment (Table 4.2.3)

Negative value of energy indicates the stability. For individual molecules, it follows: ODB > PN > MIBK > CYNO > CYN > EMK. The most stable complexes are ODB + ketones which are in the order: ODB + PN > ODB + MIBK > ODB + CYN > ODB + EMK. In Cis/Trans complex; Trans-OCP + PN is found to be most stable.

Complexes having maximum dipole moment follow: Trans-OCP + PN > Trans-OCP + EMK > Trans-OCP + MIBK.

Table 4.2.3: Optimized en	ergy and dipole moment	of systems at B3LYPLevel	using 6-31G (d) basis set.
---------------------------	------------------------	--------------------------	----------------------------

Sr. No.	Name of system	Energy (a. u.)	Dipole Moment (Debye)
1	ODB	-1151.3433	3.5027
2	EMK	-232.4678	2.8669
3	PN	-424.2079	
4	MIBK	-311.0961	2.7432
4	CYN	-235.8804	0.0002
5	Trans-OCP+EMK	-999.5437	5.7414
6	Trans-OCP+PN	-1191.2828	6.2946
7	Trans-OCP+MIBK	-1078.1724	5.4934
8	Trans-OCP+CYN	-1076.9074	5.9452

	RESEARCH JOURNEY International Multidisciplinary E-Research Journal				
(SEARCIMUUMMET	Impact Factor - (SJIF) – 6.261, Modern Trends in Green Ch	(CIF) - 3.452, (GIF) -0.676 emistry, Medicinal Chemistry and e	Special Issue - 118 nvironmental Science	February 2019	UGC Approved No. 40705
9	Cis-OCP+EMK	-999.5424		3.6160	
10	Cis OCP+PN	-1191.2734		2.7734	12.34
11	Cis- OCP+MIBK	-1078.1709		3.5196	
12	Cis-OCP+CYN	-1002.9449		0.9121	
13	ODB+EMK	-1383.9061		0.9261	
14	ODB+PN	-1575.6448		-	
15	ODB+MIBK	-1462.5344		0.8085	
15	ODB+CYN	-1387.3164		2.9289	

### 4.2.4 HOMO and LUMO

The HOMO is the orbital that could act as an electron donor, since it is the outermost (highest energy) orbital containing electrons. The LUMO is the orbital that could act as the electron acceptor, since it is the innermost (lowest energy) orbital that has room to accept electrons. Hardness is dependent on the closeness of the frontier orbitals. When the HOMO and LUMO are close together, the energy difference between the two is very low and the absolute hardness is low. In this case molecule/s can easily share orbitals to create covalent bonds/complexing. In other words, we can say strong intermolecular interactions. When the frontier orbitals are far apart, their difference in energy is large. Typically, the ionization energy in the LUMO is high in this situation. In such molecules/s, the absolute hardness is high and they are not willing to share covalent bonds i.e. weak intermolecular interactions.

Generally, polarizable molecules form strong bonds with other polarizable molecules, whereas unpolarizable molecules form strong bonds with other unpolarizable molecules.

### 4.2.4.1 HOMO:

I) Individual: for individual negative value as follows: AP > AC > CYNO > ODB > CYN.High negative value is observed in the case of AP and low value is observed in the case of CYN.

PN > MIBK > EMK > ODB > CYN.Low negative value is observed in the case of CYN.

II) Complexes: All complexes show lower negative value as compared to the corresponding individual molecules of ketones.

### 4.2.4.2 LUMO:

I) Individual: for individual negative value as follows:Negative value is low observed in the case of CYN.PN > MIBK > EMK > ODB > CYN. High negative value is observed in the case of PN and low value is observed in the case of CYN.

II) Complexes: All ODB complexes are having nearly same LUMO energy with individuals.

### Fig 4.2: Optimized geometries of molecule by DFT method atB3LYP/6-31G (d) level.



HOMO-EMK



LUMO-EMK



RESEARCH JOURNEY International Multidisciplinary E-Research Journal				
Impact Factor - (SJIF) – 6.261,	(CIF) - 3.452, (GIF) –0.676	Special Issue - 118	February 2019	UGC Approved
Modern Trends in Green Che	mistry, Medicinal Chemistry and e	Invironmental Science		No. 40705



HOMO-PN



HOMO-CIS OCP+PN



HOMO-CIS OCP+MIBK



HOMO-CIS OCP+CYN



HOMO-TRANS OCP+PN



LUMO-PN



LUMO-CIS OCP+PN



LUMO-CIS OCP+MIBK



LUMO-CIS OCP+CYN



LUMO-TRANS OCP+PN



#### HOMO-ODB+MIBK

LUMO-ODB+MIBK

#### **Conclusions:**

The compound data providence for the existence of interactions between ODB and

OCP with ketones studied by using IR Spectroscopy and theoretical Chemistry. Spectrum of some OCP + Ketones shows to absorption bands for -OH. The Sharp band is due free -OH while weak band is due to H-bonded -OH. Presence of hydrogenbonding between -OH of OCP with carbonyl oxygen of studied ketones and absence of hydrogen bonding in systems of ODB+ketones is confirmed by FTIR.Gaussian also predicts dipole moments and higher multipole moments. This term indicates the nature i.e. polar or nonpolar. The dipole moment is the first derivative of the energy with respect to an applied electric field. It is a measure of the asymmetry in the molecular charge distribution and is given a vector in three dimensions.Generally, polarisable molecules form strong bonds with other polarisable molecules, whereas unpolarisable molecules form strong bonds with other unpolarisable molecules.

### **References:**

- 1. Ezhil Pavai R, Vasantharani P & Kannappan A N, Indian J Pure Appl Phys, 2007, 45, 587.
- 2. Ali A, Yasmin A & Nain A K, Indian J Pure Appl. Phys, 2002, 40, 315.
- 3. Pimemental G C and McClellan A L; "Hydrogen Bond", W H Freeman, San Francisco (1960).
- 4. Josesten M D and Schaad L; "Hydrogen Bonding", Marcel Dekker, New York (1974).
- 5. Vinogradav S N and Linnell R H; "Hydrogen Bonding", Van NostandReinhold, New York (1971).
- 6. Legon A C and Millen D; J ChemSoc Rev, 21 (1992) 71.
- 7. Scheiner S; AccChem Res, 27 (1994) 402.
- 8. Etter M C; AccChem Res, 23 (1990) 120.
- 9. Gordon M S and Jenson J H; AccChem Res, 29 (1996) 536.
- 10. Hibbert F and Emsley J; AdvPhys Org Chem, 26 (1990) 255.
- 11. Kirby A J; AccChem Res 30 (1997) 290.
- 12. Desirajau G R; Crystal Engineering: "The design of Organic Solids", Elsevier Amsterdam (1989).

	RESEARCH JOURNEY International Multidisciplinary	ISSN- 2348-7143	
STATISTICS AND	Impact Factor - (SJIF) – 6.261, (CIF) - 3.452, (GIF) –0.676 Special Issue - 118 Modern Trends in Green Chemistry, Medicinal Chemistry and environmental Science	February 2019	UGC Approved No. 40705

13. Jeffery G A and Saenger W; "Hydrogen Bonding in BiologicalStructures", Springer-verlagBarlin, (1991).

- 14. Gerit J A, Kreevoy M M, Cleland W W and Frey P A; ChemBiol, (1997)250.
- 15. Green R D; "Hydrogen Bonding by C-H Groups", Willey: New York, 974.
- 16. Steinger T and Saenger S; J Am ChemSoc, 114 (1992) 10146.
- 17. Gu Y, Kar T and Scheiner S; J Am ChemSoc, 121 (1999) 9411.
- 18. Allerhand A and Schleyer P V R; Am ChemSoc, 85 (1963) 1715.
- 19. Turi L and Dannenberg J J; PhyChem, 99 (1995) 639.
- 20. Scheiner S, Grabowski S J and Kar T; J PhyChem, 105 (2001) 10607.
- 21. Muchall H M; J PhyChem, 105 (2001) 632.
- 22. Desiraju G R; Acc Chem Res, 29 (1996) 441.
- 23. Derewenda Z S, Lee L and Derewenda U; J Mol Bio, 252 (1995) 248.
- 24. Metzger J and Pullman A; Bull SocChim France, 1166 (1948).
- 25. Zahradnik R and Kontecki J; Coll Czech ChemComn, 26 (1961) 156.
- 26. Vitry J and Metzger J; Bull SocChim France, 1784 (1963).
- 27. Bansal R K, Karaghiosoff K, Gupta N, Schmidpeter A and Spindler C; ChemBer, 124 (1991) 475.
- 28. Bansal R K, Gupta N, Kabra V, Spindler C, Karaghiosoff K and Schmidpeter A; Heteroatom Chem, 3 (1992)259.
- 29. Gupta N, Jain C B, Heinicke J, Bharatiya N, Bansal RK and Jones P G;Heteroatom Chem,9(1998) 259.
- 30. Wang B, Hinton J F and Polay P; J PhyChem, 107 (2003) 4683.
- 31. Cubero E, Orozco M, Hobza P and Luque F J; PhysChem, A 103(1999) 6394.
- 32. Hobza P and Havlas Z; ChemPhysLett, 303 (1999) 447.
- 33. Scheiner S and Kr T; J PhyChem, A 106 (2002) 1784.
- 34. Kadam U B, Hasan Mehdi and Sawant A B, International Journal of Chemical Sciences 4 (2) (2006) 458.
- 35. Nazir H, Yildiz M, Yilmaz H, Tahir M N and Ulku D; J MolStruct, 524 (2000) 241.
- 36. Cuma M, Scheiner S and Kar T; J Am ChemSoc, 120 (1998) 10497.
- 37. Filarowski A, Glowiak T and Koll A; J MolStruct, 484 (1999) 75.
- 38. Raissi H, Moshfeghi E, Jalbout A F, Hosseini M S and Fazli M; Int J Quant. Chem. 107 (2007) 1835.
- 39. Cuma M, Scheiner S and Kar T; J MolStruct, (Theochem) 467 (1999) 37.
- 40. Chung G, Kwon O and Kwon Y; J PhysChem, A 102 (1998) 2381.
- 41. Fores M and Scheiner S; ChemPhys 246 (1999) 65.
- 42. Cheng C, Shyu S F and Hsu F S; Int J Quant Chem, 74 (1999) 395.
- 43. Barone V, Palma P and Sanna N; PhysLett, 381 (2003) 45.
- 44. R M Silverstein and F X Webster; "Spectrometric Identification of Organic Compounds"

6<sup>th</sup>ed New York (2006).

45. Frisch M J, Trucks G W, Schlegel H B, Scuseria G E, Gill P M W, Johnson B, Chen W, Wong M W, Gonzalez C and Pople J A; Gaussian, Inc., Wallingford CT, 2004.

cipal

Kr. V. N. Nalk Shikehan Prasarak Sanstha's Arts, Commerce & Science College, Canada Cernar, Nashik.